

## TRAUMATIC POSTERIOR HIP DISLOCATION IN 6 YEAR OLD CHILD: A CASE REPORT

Ahmed Parvez<sup>1</sup>, Tomar Sagar<sup>2</sup>, Menwal Gaurav<sup>3</sup>, Verma Jaiprakash<sup>4</sup>, Devberma Ijack<sup>5</sup>

<sup>1</sup>Professor, <sup>2,4,5</sup>Junior Resident, <sup>3</sup>Senior Resident,  
IIRM Medical College, Meerut India

**Corresponding Author:**

E-mail: drahmedparvez@yahoo.com

### INTRODUCTION

A review of the literature confirms that traumatic dislocation of the hip in children is rare. The aim of this study is to report the case of a posterior hip dislocation in a 6 year old child due to fall while playing. The diagnosis of dislocation in children is difficult and such trauma can lead to serious complications.

### CASE REPORT


A case of 6 year old girl was admitted to emergency department of S.V.B.P. Hospital, Meerut, with chief complaints of pain, swelling and deformity in right hip region for 3 days. The trauma was caused while playing when another child pushed her from behind.

On clinical examination, there was pain, restricted range of motion and the attitude of right hip was in slight flexion, adduction and internal rotation. Maximal discomfort was localized to the right buttock, which was tensely swollen, tender and warm to the touch. The examinations done on other joints did not reveal any abnormality regarding the range of motion or ligament laxity. A plain x-ray was done revealing posterior and superior displacement of the head of the femur out of the acetabulum without any accompanied fracture. There was no evidence of infective pathology on routine investigations.



**Fig 1: X-ray showing posterior dislocation of (Rt) Hip AP view (A) lateral view (B)**

Under general anesthesia, the hip was reduced, using closed method. The hip joint was stable in full range of motion. The hip was kept in below knee fixed skin traction on TK splint. Post-reduction radiographs showed a concentrically reduced hip without any associated fracture or asymmetry of joint space.



**Fig 2: X ray showing concentrically reduced (r) hip joint with appl TRACTIO traction**

After 2 days of skin traction, the patient was discharged from the hospital with advice not to bear weight on the affected limb for 6 weeks. At 6 weeks post-reduction, clinical and radiological assessments revealed no abnormality.

## DISCUSSION

A review of the literature confirms that traumatic dislocation of the hip in children is indeed rare<sup>1,2,3</sup> and is frequently associated with trivial degrees of trauma, especially in children under the age of 5-6 years.<sup>2,4,5</sup> The diagnosis is usually made promptly by recognition of the classic deformity, i.e. shortening with adduction, flexion and internal rotation, but there are many reported cases of delayed diagnosis.<sup>1-3,6-10</sup> The most potent cause of a missed diagnosis appears to be associated femoral shaft fracture<sup>6,9</sup>, the fracture directs attention away from the hip and may obscure the usual deformity by its own displacement. Other reasons for missing the diagnosis are (as in this case) the misleading minor degree of trauma and the rarity of the condition. Multiple injuries may also obscure the diagnosis.<sup>6,10</sup> It is mollifying to note that the literature suggests that in cases of ipsilateral femoral shaft fracture the diagnosis was missed less often in the pre-radiography era than it is now. Anterior traumatic dislocation of the hip is far less frequent than the posterior form<sup>1-3</sup> and the diagnosis is usually obvious. The chief associated problem is compression of the femoral vessels. Early diagnosis and reduction of a dislocated hip is essential to prevent its chief complication, avascular necrosis of the femoral head, the overall incidence of which is between 4% and 10%.<sup>3,7,10,11</sup> It is universally agreed that early reduction (within 24 hours) and the age of the patient (less than 5 years) are major factors in reducing this incidence.<sup>2,3,12</sup> Most authors state that a severe degree of initial trauma predisposes to avascular necrosis,<sup>11,12</sup> but some disagree.<sup>8,13</sup> Sciatic nerve damage<sup>4,6,9,12</sup> and associated acetabular fractures<sup>2</sup> are very rarely reported. Other uncommon complications of traumatic hip dislocation in children are late osteoarthritis, with or without coxa magna<sup>7,10</sup> and recurrent dislocation.<sup>5,14</sup> The treatment of traumatic dislocation of the hip is prompt reduction by closed manipulation performed under general or spinal anaesthesia. This is usually relatively easy. Rare cases of failed manipulation are either due to buttonholing of the femoral head through the hip capsule or infolding of the labrum acetabulare, and in such cases open reduction is necessary. There is no uniformity of opinion about the length of time required for immobilization, but a distillation of the literature would suggest that 6 weeks in traction or hip spica followed by immediate weight-bearing is reasonable. There is, however, no specific evidence to support this regimen.

## REFERENCES

1. Bunnel WP, Webster DA. Late reduction of bilateral traumatic hip dislocation in a child. *Clin Orthop* 1980; 147: 160-163.
2. Funk FJ. Traumatic dislocation of the hip in children. *Bone Joint Surg (Am)* 1962; 44: 1135-1145
3. Pennsylvania Orthopaedic Society. Traumatic dislocation of the hip joint in children. *J Bone Joint Surg (Am)* 1968; SO: 79-88.
3. Schlonsky J, Miller PR. Traumatic hip dislocations in children. *J Bone Joint Surg* 1973; 55: 1057-1063.
4. Simmons RL, Elder JD. Recurrent post-traumatic dislocation of the hip in children. *South Med J* 1972; 65: 1463-1466.
5. Debn E, Immernan EW. Dislocation of the hip combined with fracture of the shaft on the same side. *J Bone Joint Surg* 1951; 33: 731-745.
6. Glass A, Powell HDW. Traumatic dislocation in the hip of children. *J Bone Joint Surg* 1961; 43: 29-37.
7. Halibunon RA, Brokenshire FA, Barber JR. Avascular necrosis of the capital femoral epiphysis after traumatic dislocation of the hip in children. *J Bone Joint Surg* 1961; 43: 43-46.
8. Helal B, Skevis X. Unrecognised dislocation of the hip in fracture of the femoral shaft. *J Bone Joint Surg (Br)* 1967; 49: 293-300.
9. MacFarlane I, King D. Traumatic dislocation of the hip joint in children. *Aust NZ J Surg* 1976; 46: 227-231.
10. Piggot J. Traumatic dislocation of the hip in children. *J Bone Joint Surg (Br)* 1961; 43: 38-42.
11. Barquet A. Traumatic hip dislocation in childhood. *Acta Orthop Scand* 1979; SO: 549-553.
12. Freeman GE. Traumatic dislocation of the hip in children. *J Bone Joint Surg* 1961; 43: 401-406.
13. Liebenberg F, Dommissie GF. Recurrent post-traumatic dislocation of the hip. *J Bone Joint Surg* 1969; 51: 632-637.
14. Fordyce AJW. Open reduction of traumatic dislocation of the hip in a child. *Br J Surg* 1971; 58: 705-707.
15. Pennsylvania Orthopaedic Society. Traumatic dislocation of the hip joint in children. *J Bone Joint Surg* 42-A : 705- 710.
16. Piggot J. Traumatic dislocation of the hip in children. *J Bone Joint Surg* 1961 ; 43-B : 38-42.
17. Rieger H, Pennig D, Klein W, Grünert J. Traumatic dislocation of the hip in young children. *Arch Orthop Trauma Surg* 1991 ; 110 : 114-117.
18. Sahin V, Karakas ES, Turk CY. Bilateral traumatic hip dislocation in a child : a case report and review of the literature. *J Trauma* 1999 ; 46 : 500-504.
19. Schlonsky J, Miller PR. Traumatic hip dislocations in children. *J Bone Joint Surg* 1973 ; 55-A : 1057-1063.
20. Loupasis G, Morris EW. Asymmetric bilateral traumatic hip dislocation. *Arch Orthop Trauma Surg* 1998 ; 118 : 179-180. Macfarlane I, King D. Traumatic dislocation of the hip joint in children. *Aust N Z J Surg* 1976 ; 46 : 227-231.
21. Mehlman CT, Hubbard GW, Crawford AH, Roy DR, Wall EJ. Traumatic hip dislocation in children. Long term follow-up of 42 patients. *Clin Orthop* 2000 ; 376 : 68-79.
22. Muratli HH, Dagli C, Bicimoglu A, Tabak AY. Recurrent traumatic hip dislocation in a child : Case report. *Acta Orthop Traumatol Turc* 2004 ; 38 : 149-153.
23. Nirmal Kumar J, Hazra S, Yun HH. Redislocation after treatment of traumatic dislocation of the hip in children

- : a report of two cases and literature review. Arch Orthop Trauma Surg 2008 Aug 22.
24. Offierski CM. Traumatic dislocation of the hip in children. J Bone Joint Surg 1981 ; 63 : 194-197.
  25. Pearson DE, Mann RJ. Traumatic hip dislocation in children. Clin Orthop 1973 ; 92 : 189-194.
  26. Simmons RL, Elder JD. Recurrent post-traumatic dislocation of the hip in children. South Med J 1972 ; 65 : 1463- 1466.
  27. Vialle R, Odent T, Pannier S et al. Traumatic hip dislocation in children. J Pediatr Orthop 2005 ; 25 : 138-144.